

INORGANIC CHEMISTRY

Lesson 9

Salts and their reactions.

Reaction of active metals with water.

Bases, basic oxides, and their interaction with acids.

November 12, 2017

1 Nomenclature of salts

Before we started to discuss reactions of salts, and methods of their preparation, we need to know how to name them. Since salt are composed of some acidic residue¹ and a metal atom (or atoms), it would be logical to derive salt's name from the names of corresponding acids and metals. That is exactly what chemists do. The salt nomenclature (naming rules) are simple and straightforward. The rules are as follows.

Rule 1. (for non-oxygen acids):

The name of a salt formed by some metal and some non-oxygen acid is formed by combining the metal's name and the name of the acid. The prefix 'hydro' is removed, and the ending 'ic' is replaced with 'ide'

Examples Hydrochloric (*hydro-chlor-ic*) acid (HCl) and sodium form a salt named 'sodium chloride (NaCl).

Calcium and hydrochloric acid form calcium chloride CaCl_2 .²

Rule 2. (for oxygen containing acids)

In salts formed by 'ic' acids, the ending 'ic' is replaced with 'ate'.

Examples Carbonic (carbon-*ic*) acid, H_2CO_3 and sodium form a salt named 'sodium carbonate (Na_2CO_3).

Sulfuric acid and calcium form a salt named 'calcium sulfate (CaSO_4).

Rule 2a. (for 'ous' acids).

¹Acidic residue is what is remaining when one or several hydrogens have been removed from the acid's molecule.

²In school, you may be requested to include numerals into the salt's name. For example, instead of 'calcium chloride' (CaCl_2) you should say 'calcium *dichloride*', instead of 'aluminum chloride' (AlCl_3) you should say 'aluminum *trichloride*', etc. Although that is not incorrect, chemists prefer not to do that. During our lessons we will not use numerals unless that is absolutely necessary.

In salts formed by 'ous' acids, the ending 'ous' is replaced with 'ite'.

Examples. Sulfur-ous acid (H_2SO_3) and sodium from a salt named 'sodium sulfite' (Na_2SO_3).

Nitrous acid (HNO_2) and potassium from a salt named potassium nitrite (KNO_2).

Since the acids whose name starts with 'hypo' or 'per' also can form salts, a separate rule exists for that type acids. This rule is simple.

Rule 3.(for the acids whose names begin with 'per' or 'hypo'.)

If acid's name starts with the prefixes 'per' or 'hypo', such a prefix is preserved in the salt's name.

Examples. A salt formed by sodium and hypochlorous acid (HClO) acid is called 'sodium hypochlorite' (NaClO). A salt formed by sodium and perchloric acid (HClO_4) is called 'sodium perchlorate' (NaClO_4).

We are almost finished with the rules. There is one more rule; it deals with a situation when a metal can exist in more than one valence state. For example, iron can be either di- or trivalent, and, accordingly, it can form two different salts with hydrochloric acid. Their formulas are FeCl_2 and FeCl_3 . As we can see, both these salts are 'chlorides'. Two alternative ways exist to create unique names for each of them. Firstly, we can show metal's valence using a Roman number.

Rule 4a.

When more than one valence states is possible for some metal, its valence can be indicated with a Roman letter after the metal name.

Examples. The name of FeCl_2 is 'iron (II) chloride'. The name of FeCl_3 is 'iron (III) chloride'.

Another possibility is to add the ending 'ous' or 'ic' to the *Latin* name of the metal. The meaning of these prefixes is the same as for acids: lower valence is denoted by 'ous', whereas higher valence by 'ic'.

Rule 4b.

When some metal is in a lower valence state, that fact can be indicated by adding the ending 'ous' to its Latin name. For metals in a higher valence state, the ending 'ic' is used.

Examples. A Latin name of iron is 'ferrum', therefore FeCl_2 is called 'ferrous chloride', and FeCl_3 is called 'ferric chloride'. In Latin, copper's is called 'cuprum', therefore, their salts have names 'cuprous chloride' (CuCl), and 'cupric chloride' (CuCl_2).

This is an almost exhaustive set of rules. Most salts are named according to them. I neither expect nor want you to memorize all these rules, I presented them just to demonstrate that the rules are relatively simple. However, in future, you will use these rules frequently, and you know where can you find them: they will always be at our school's web site. The most frequently used rules are Rule 1 and Rule 2, sometimes, a Rule 4a. We will need them right now, during this lesson.

2 Reactions of salts with metals. More on the reactivity series.

Experiment 15a

Pour 5 mL of the solution of copper (II) sulfate (CuSO_4) into a test tube, and immerse an iron rod into it. Describe your observations.

After 5 minutes, the rod becomes reddish; the red film become more and more thick, and finally it peels off like a red sponge. It is easy to demonstrate that this is a copper metal. If we wait long enough we can see that a blue color of the copper sulfate solution gradually disappears, and the solution becomes pale greenish. This color belongs to another salt, iron (II) sulfate. Obviously, a chemical reaction occurs, and we can write the following equation for it:



As in the case of iron and sulfuric acid, a substitution reaction occurs, however, instead of elementary hydrogen, another simple substance, copper, is formed. Is such a reaction common for an arbitrary salt-metal pair? Let's try to think logically. Since a chemical reaction will hardly go into the opposite directions under the same conditions, swapping the sides in the equation 1 would lead to the equation describing the reaction that doesn't go normally. Actually, this consideration is totally correct: it is easy to demonstrate that reaction is not possible.

Experiment 15b


Pour 5 mL of the solution of iron (II) sulfate (FeSO_4) into a test tube, and immerse a piece of copper wire into it. Do you see any reaction?

As we expected, no reaction takes place. Maybe, this reaction is specific to copper sulfate only? Let's check that.

Experiment 15b

Pour 5 mL of the solution of silver nitrate (AgNO_3), and immerse a piece of copper wire into it. What do you see?

We see that lustrous crystals quickly form on the copper surface, and that the colorless solution becomes green-blue. It is easy to demonstrate that the crystals we obtained are a silver metal. With regard to the color, it is characteristic for most copper salts, so we can conclude some copper salt, is formed. Indeed, a chemical analysis can demonstrate a new salt forms, a copper nitrate. The equation of this reaction is as follows:



Again, similar to the of iron and copper sulfate case, a new metal and a new salt form. It is easy to demonstrate that the opposite reaction wouldn't go.

Do those reaction have anything in common? Yes. First of all, these reactions are *substitution reactions*. Secondly, in each of those reaction, one metal substitute another in its salt. Thirdly, there reactions go in one direction (copper sulfate reacts with iron), and do not go in another (iron sulfate does not react with copper). It is also easy to demonstrate that the acidic residue does not affect this reaction: for example, copper chloride and copper sulfate react similarly. All of that can be summarized as follows: copper *salts* react with iron to produce copper *metal*, silver *salts* react with copper to produce silver *metal*. It can be demonstrated that other metals, such as zinc, tin, etc, produce silver in the reaction with silver nitrate, but gold or platinum do not react with AgNO_3 . Can we summarize all these facts in some simpler rule? We can try. To do that, let's look at the reactivity series again.

Reactivity series of metals

K, Na, Li, Ca, Mg, Al, Ti, Mn, Zn, Cr, **Fe**, Co, Ni, Sn, Pb (H) **Cu**, W, Hg, **Ag**, Au, Pt

As we can see, the metals we experimented with are arranged in the following order (in terms of their reactivity): iron is left of copper, and copper is left of silver. It can be easily demonstrated that rule is general: all metals left of tin are able to substitute it from tin salts, all metals left of gold can substitute it from gold salts, and so on. This is a general rule the reactivity series was built on. And it can be summarized as follows:

More active metals substitute less active metals from their salts. The opposite reaction does not go.

3 Reactivity series and hydrogen

Let's look again at these two equations:



and



Clearly, they have much in common: both of them are substitution reactions, in both of them one element substitutes another element from its compound. The only difference is that in the first case iron reacts with salt, and in the second case it reacts with acid. This comparison explains why did we put hydrogen into the reactivity series: the metals left of hydrogen are more active than hydrogen, and they substitute hydrogen from acids (in the same manner they substitute, for example, copper or silver from their salts). In contrast, the metals left to hydrogen are less active, so they cannot substitute it from acids.

The reactivity series is a first example of an empiric law that allows us not only to describe chemical reactions, but also to predict their outcome. There are many similar law in chemistry that make it not just a random collection of facts, but a systematic science. We will learn many of them during this and the next year.

4 On the very left part of the reactivity series.

As we already know, all metals left of hydrogen are able to react with acids to produce hydrogen. The closer to the left end of the series they are, the more actively they react with acids. Thus, almost no reaction takes place between acetic acid and iron; zinc does react with acetic acid, although very slowly; magnesium reacts with acetic acid very actively. In other words, active metals are capable of substituting hydrogen even from the molecules that are not willing to easily donate hydrogen. What about more active metals, the metals left of magnesium? Those metals (shown in bold below) are so active that they react too violently even with acetic acid.

Reactivity series of metals

K, Na, Li, Ca, Mg, Al, Ti, Mn, Zn, Cr, Fe, Co, Ni, Sn, Pb (H) Cu, W, Hg, Ag, Au, Pt

We know water contains hydrogen, but its hydrogen is not too active (at least, no hydrogen is formed when water is added to zinc or iron). We also know that common metals are not capable of substituting hydrogen from water. Let's see if active metals can do that.

Experiment 16

Pour 100 mL of water into a glass beaker. Take approximately 1/4 gram of sodium metal, carefully drop it into the beaker, and immediately cover the beaker with a glass funnel. Attach a test tube to the narrow end of the funnel as shown on the Fig. 1. When the reaction is complete, remove the test tube (do not flip it) and bring a burning candle to the tube's neck. Describe your observations.

We can see sodium's behavior differs dramatically from what we expect from a metal. It reacts with ordinary water, a compound we usually consider inert (not active) in normal conditions, and some gas is formed during that reaction. The reaction is so violent, that the sodium melts and collapses into a lustrous ball on the water surface.³ As we demonstrated, the gas formed during this reaction is combustible,⁴ and it is lighter than air (it goes up, and it stays in the test tube when its neck is pointing down). We will see later that are the properties of hydrogen. We can conclude therefore that reaction of sodium with water produces the same gas we obtained during the reaction between zinc and hydrochloric acid, i.e. hydrogen. Had we tried to evaporate the remaining liquid in the beaker,⁵ we would



Figure 1: Reaction of sodium with water. A liquid sodium ball in the left is partially covered with a water mist.

³Density of sodium is smaller than the density of water, so sodium does not sink in water.

⁴Sometimes, sodium becomes so hot during this reaction that the gas ignites spontaneously.

⁵We will not do that for safety reasons.


obtain a white soapy substance with a formula ‘NaOH’. Therefore, the equation of this reaction can be written as follows:




In this equation, one compound (NaOH) has a formula we have never seen before. It is not an oxide, not a salt, and not an acid. We have never dealt with a compound of such a type before. Definitely, NaOH belongs to a new type of compounds.

5 Hydroxides, bases and alkali.

A compound with a formula NaOH can be represented as an analog of sodium oxide Na_2O , where one sodium atom has been replaced with a hydrogen atom:



Sodium oxide



Sodium *hydroxide*

It is intuitively clear why we call it “hydroxide”: in this substance, one oxygen’s valence is occupied by a hydrogen atom, and this diatomic particle, which is called a “hydroxy group”, forms a compound with the atom of metal (in this concrete case, sodium).

Hydroxides are the compounds with general formula $\text{M}(\text{OH})_n$, where M denotes some element, and ‘n’ (a number of hydroxy groups) is equal to the element’s valence. Hydroxides, form another major class of inorganic compounds.

Examples of hydroxides are calcium hydroxide $\text{Ca}(\text{OH})_2$, magnesium hydroxide $\text{Mg}(\text{OH})_2$, aluminum hydroxide $\text{Al}(\text{OH})_3$.


5.1 Nomenclature of hydroxides

In contrast to acids, there is almost nothing you need to remember about names of hydroxides. The hydroxide’s name is created by combining the element’s name and the word “hydroxide”, in the same way we did in the end of the previous section. A situation is a little bit more complicated for the elements that can exist in more than one valence state. As a rule, such elements can form more than one hydroxide. In that case, you must denote the element’s valence with a Roman numeral. For example iron forms two hydroxides: iron (II) hydroxide ($\text{Fe}(\text{OH})_2$), and iron (III) hydroxide ($\text{Fe}(\text{OH})_3$). That’s all what we need to know about hydroxides naming rules.

5.2 Preparation of hydroxides. Part I.

Our observations we made during the Experiment 16 can be generalized: it is natural to expect other metals in the left part of the reactivity series are able to react with water. This

our hypothesis is correct, and it is easy to prove experimentally that potassium, lithium, calcium, as well as some other active metals not included in the reactivity series⁶ produce hydrogen and hydroxides during their reaction with water according to the scheme:



where Me denotes some active metal, and ‘n’ is the metal’s valence. **Reaction of active metals with water is the first method to obtain hydroxides.** Is it the only method? Apparently, no. Alternative ways to prepare of hydroxides exist.

Experiment 17

Place approximately 10-20 grams of calcium oxide⁷ into a glass beaker or a steel bowl. Add approximately 10 mL of water to it. What happens to the solid? Carefully touch an external surface of the beaker. What do you feel?

We can see that during this experiment a thin white powder of calcium oxide absorbs water and swells. The volume of the solid increases, and large amount of heat evolves. The reaction mixture becomes so hot that water we added starts to boil, and a cloud of steam forms above the beaker. The compound that forms during this reaction is calcium hydroxide ($\text{Ca}(\text{OH})_2$). Since ancient times, it is being used to prepare a mortar, or a lime slurry. Until XIX century, when modern cement had been invented, slaked lime was vital and indispensable for construction. The equation of this reaction is:



Due to its practical importance, this reaction has its own name: “slaking”,⁸ and, accordingly, the technical name of calcium hydroxide is “slaked lime”.

Ancient people started to use slaked lime long before a concrete had been discovered. They used it to prepare lime mortar, a paste to bind bricks or stone construction blocks together. During several months after the construction ended, the binding mixture was becoming harder and harder, and eventually it converted into some artificial stone. An astonishing longevity of ancient Roman aqueducts and bridges (for example, the aqueduct at the figure 2), which had been built more than two thousand years ago, and, nevertheless, remain virtually unchanged, may serve as a perfect demonstration of an excellent quality of the lime mortar cement. During this and the next lesson we will learn about the chemistry behind that phenomenon.

⁶Remember, we are dealing with a short version the reactivity series: some other metals, besides potassium, sodium, lithium and calcium, are capable of reacting with water to produce hydrogen and hydroxides.

⁷Calcium oxide (CaO) is known under a name “quick lime”.

⁸This term refers only to the reaction of calcium oxide and water. No special term exist for the reaction of other oxides.

Calcium oxide is not the only oxide capable of formation of hydroxides. Many other oxides, such as sodium oxide or potassium oxide, react with water. The equation of the reaction of sodium oxide and water is as follows:



The reaction between metal oxides and water is the second way to prepare hydroxides.

6 Soluble hydroxides and their properties. Bases. Alkali.

Now it is a time to look at the properties of the compounds we prepared during the experiments 16 and 17.

6.1 Sodium hydroxide.

Sodium hydroxide (NaOH), also known as lye is a white highly hygroscopic solid with a slippery feeling upon contact.⁹ It is highly soluble in water; its dissolution is a highly exothermic process. Accordingly, sodium hydroxide belongs to a family of compounds called “soluble hydroxides”. Another common name for soluble hydroxides is “alkali”.

Soluble metal hydroxides are called “alkali”.

A solution of NaOH corrodes skin, paper, fabric, and many other materials changes the color of a standard indicator paper to deep blue, which corresponds to pH values above 13. Addition of few drops of the phenolphthalein¹⁰ solution produces a purple color. This is indication that sodium hydroxide is *basic*. Basicity is a common property of soluble hydroxides, such as potassium hydroxide (KOH) and lithium hydroxide (LiOH).

6.2 Calcium hydroxide.

In contrast to sodium hydroxide, calcium hydroxide is poorly soluble in water. However, its water solution, a so called “lime water”,¹¹ is also basic, because it produces a pink color when a colorless solution of phenolphthalein is added to it. Other moderately soluble hydroxides, such as silver hydroxide (AgOH) have similar properties.

⁹Actually, you should never touch it! It is very corrosive, so in a case of contact with your skin you must immediately rinse it with large amount of water until a slippery feeling disappears.

¹⁰An indicator. This substance is colorless, but it becomes purple when a common pH paper becomes deep-blue.

¹¹We are already familiar with it. Do you remember what we used it for?

6.3 Properties of soluble hydroxides. Reaction with acids (“Neutralization”).

Thus we have learned that the alkali are “basic”. We also know acids are “acidic”. So far, we do not fully understand what does it mean, however, we know that a standard indicator paper becomes red in an acid media, and it becomes deep-blue in a basic media. We also know that an indicator dye, phenolphthalein, becomes pink in a presence of a base. In connection to that, it is interesting to see what happens when an acid and a base are mixed together.

Experiment 18.

Pour 100 mL of 4% solution of sodium hydroxide into a glass beaker and add few drop of phenolphthalein solution to it. Using a pipette, gradually add a 3.7 % dilute solution of hydrochloric acid to the beaker. When a pink color disappeared, record the volume of the HCl solution you added. Pour the solution obtained into a porcelain bowl and evaporate on open fire. What did you obtain?

Interestingly, during this experiment, exactly 100 mL of the HCl solution are needed for the pink color to disappear. What conclusion can be drawn from that fact? You have probably noticed that the concentration of both solutions was chosen in such a way that equal volumes of NaOH and HCl solutions contained equal number of NaOH and HCl molecules. Indeed, the molecular weight of NaOH is 40 Da ($23 + 16 + 1 = 40$), and the molecular weight of HCl is approximately 37 ($35.5 + 1 = 36.5$). That means, 100 mL of 4% NaOH solution contain 4 g of NaOH; the same volume of 3.7% HCl solution contains 3.7 g of HCl, and these amounts are proportional to the molecular masses of NaOH and HCl, accordingly. In other words, by the end of our experiment, one HCl molecule have been added per one NaOH molecule. What we obtained as a result of that addition? Clearly, the solution is not basic any more: a pink color of phenolphthalein had disappeared. Is it acidic? No. An indicator paper immersed into the solution does not become red.

It is easy to demonstrate that, when the order of addition changes (in other words, if 100 mL of a 4% NaOH solution will be added to 100 mL of 3.7% HCl solution), the result will be the same.

That means an acid and a base, when mixed together, “destroy”, or “eliminate” each other: an acid eliminates “basicity”, and a base eliminates “acidity”. Such a mutual elimination is called “neutralization”.

An acid and a base react with each other. Such a reaction is called “neutralization”. The resulting solution is neither acidic nor basic; it is “neutral”.

What is a product of the neutralization reaction? Evaporation of the solution obtained in the Experiment 18 yields white crystals having a cubic shape. The shape of the crystals, their density, hardness, melting temperature, and other physical properties¹² are totally identical


¹²Of course, we have neither an opportunity nor time to do all those measurement in the class, but, have we done that, we would obtain exactly the same results.

to the properties of common salt. Therefore, we can conclude the product of this reaction is sodium chloride (NaCl), and the reaction's equation is as follows:



(The fact that the second product of this reaction is water also can be confirmed experimentally. However, we are not doing that because of the lack of time.)

Other hydroxides react with hydrochloric acid similarly. For example, calcium hydroxide produces calcium chloride (CaCl_2):



Other acids are also react with the bases in the same way. Thus, potassium sulfate is formed in the reaction between potassium hydroxide and sulfuric acid:



Numerous experiments demonstrated that every acid and every hydroxide can react with each other to produce salt and water, and this is a fundamental property of every base and every acid.¹³ These observations had been summarized by a Swedish chemist Svante Arrhenius, who proposed a first definition of the term “bases”.

Bases are the compounds that are capable of donating a hydroxy group in reaction with acids. Acids are the compounds donating hydrogen in a reaction with bases. The products of the reaction between an acid and a base (a neutralization reaction) are salt and water.

Historically, this is a first definition of the terms “acid”, “base” and “salt”. Current definitions of the terms “acid” and “base” are somewhat broader. However, since these old definitions work perfectly for our purposes, we will stick with them during this school year.



Figure 2: Svante Arrhenius (1859-1927)

A man who proposed the first definition of “acids” and “bases”.

6.4 Reaction of bases with acidic oxides.

If we pour transparent lime water in a glass beaker and blow bubbles through it, the water will become turbid in few minutes. What happens during this reaction? In this reaction, carbon dioxide you exhale react with calcium hydroxide according to the equation:



¹³Of course, when the acid is too weak and the hydroxide is poorly soluble, such a reaction would be too slow to be detected.

The salt that forms during this reaction is calcium carbonate, aka calk, or limestone, of marble. Since this compound is insoluble in water, even small amount of CO_2 are sufficient to turn a clear and transparent lime water (i.e. $\text{Ca}(\text{OH})_2$ solution into a turbid suspension. More importantly, a solid $\text{Ca}(\text{OH})_2$ is also capable of reacting with atmospheric CO_2 , which leads to a gradual conversion of soft, semi-soluble slaked lime ($\text{Ca}(\text{OH})_2$) into a rock-solid limestone (CaCO_3).

As we can see, in this reaction, a salt, is formed not in a reaction between an acid and a base, but in the reaction between an *acidic oxide* and a *base*.

Again, that is a common property of all bases:

Bases are capable of reacting with both acids and acidic oxides to form salts. The same salt is formed in the reaction with an acid and a parent acidic oxide.

A question. Do you understand now the chemical mechanism of the astonishing longevity of Antique Roman aqueducts, bridges and buildings?

7 Bases and basic oxides.

From the section 2.2, we know that some metal oxides are capable of reacting with water, and the metal hydroxide is a major product of such a reaction. In connection to that, it is interesting to compare that reaction with the reaction we studied during the Lesson 6 (section 2). As we already know, the product of the reaction between phosphorus (V) oxide and water is phosphoric acid:



Let's compare it with a reaction between calcium oxide and water (equation 7). Since phosphorus (V) oxide produces an acid, we call it, as well as all other oxides of that type, an "acidic oxide". Accordingly, it would be logical to introduce a term "basic oxide" to denote an oxide that produces a base in reaction with water.

Basic oxides are the oxides that react with water to produce a base.

7.1 Reaction of basic oxides with acids.

In the previous section, we found that acids and acidic oxides yield the same product in a reaction with acids. What about the bases and the basic oxides?

Experiment 19.

Put 1-2 grams of calcium oxide into a glass beaker. Add 10 mL of dilute acetic acid to it. When the reaction stopped, put the beaker on a hot plate and gently evaporate to dryness.¹⁴ What did you obtain?

¹⁴Actually, we will not do this experiment, because it should be done either in the fume hood or on open air: although the vapors of acetic acid are not toxic, they have a very unpleasant odor.

The solid we obtained is a salt called “sodium acetate”. It is the same salt you would obtain, had you taken calcium hydroxide instead of calcium oxide. Again, we can experiment with a various basic oxides and acids, and the result will be the same:

Acids are capable of reacting with both bases and basic oxides to form salts. The same salt is formed in the reaction with an base and a parent basic oxide.

You have probably noticed that the above statement mirrors the statement from the section 3.4: the only difference between them is that the words “acid” and “base” have been swapped.¹⁵ Such a symmetry is not merely a coincidence. Let’s talk about that during the next lesson.

Homework

1. A tin-lead alloy or silver can be used as a solder to connect copper parts together. Imagine you have two copper vessels. One of them was made using a tin-lead solder. Silver was used to solder the parts of another vessels. Can these vessels be used to store dilute hydrochloric acid?
2. Brass is a copper and zinc alloy. Depending on a zinc/copper ratio, it may have different properties. You have brass shavings with unknown zinc content. You took 5 grams of those shavings and added 200 mL of 20% HCl¹⁶ to them. You used an apparatus (similar to that we used for preparation of oxygen) that allowed you to collect all gas that forms during that reaction. When the reaction was complete (evolution of the gas ceased) you measured the gas volume, and found that 0.4 L of some gas was formed. Can you determine a zinc content (in percents) in the brass using these data? For your calculations, you can use the fact that at room temperature and under atmospheric pressure, two grams of hydrogen occupy 22.4 L.¹⁷
3. It is a good time to start summarizing what we learned during previous lessons. Please, answer, how many major classes of inorganic compounds have we currently learned? Can you list them, and briefly describe their properties?
4. Since we are starting to discuss properties of various chemical compounds (we already know almost all major classes of inorganic compounds), we need to practice in naming them. Please, name the following compounds:
 - a. $\text{Al}(\text{OH})_3$
 - b. FeSO_3
 - c. H_3PO_4
 - d. MgCO_3

¹⁵That is what I actually did: I used copy-past to save time.


¹⁶‘20%’ means that 100 mL of solution contains 20 grams of HCl.

¹⁷Generally speaking, if a mass of one molecule of some gas is X Da, then X *grams* of this gas will occupy a volume of 22.4 L at room temperature and atmospheric pressure. This is a general law named ‘Avogadro’s’ law (after its discoverer, Amedeo Avogadro).

- e. CuOH
 - f. Al₂O₃
 - g. Sn₃(PO₄)₂
 - h. K₂SO₃
5. Write formulas and name the compounds formed during the reaction between:
- a. Silver hydroxide and nitric acid.
 - b. Calcium and oxygen.
 - c. Sodium hydroxide and sulfur (VI) oxide.
 - d. Magnesium and hydrochloric acid.
 - e. Iron (II) oxide and hydrochloric acid.
 - f. Carbon and oxygen.
 - g. Iron and dilute sulfuric acid.¹⁸
 - h. Silicon oxide and potassium hydroxide.

6. Propose a reaction scheme (draw each equation separately) for the following transformation:

a.



b.



c.



d.



7. Propose three different reactions to prepare each of the following compounds:
- a. Calcium chloride.
 - b. Iron (II) sulfate.
 - c. Aluminum nitrate.
8. Propose four different reactions to prepare sodium phosphate Na₃PO₄.
9. You took 20 grams of calcium oxide, added 300 mL of 10% HCl, and evaporated the solution obtained to dryness. What is the formula of the solid you obtained? What is its mass?
10. A piece of zinc reacted with dilute hydrochloric acid in the apparatus that allowed you to collect all the gas formed. The zinc dissolved completely, and the volume of gas

¹⁸When polyvalent metals, such as iron, react with acids, a salt is formed where the metal has the lowest possible valence.

you collected was 35L. What is the amount of the second product formed during this reaction? What was the amount of zinc taken?¹⁹

11. Pont du Gard aqueduct was built 2000 years ago. It is still rock-stable. What chemical reaction is behind its outstanding stability? (A hint: we discussed this reaction in the class).

As usual, I would be grateful if you sent me your homework by evening of next Saturday. My e-mail is mark.lukin@gmail.com.

If you want to draw structural formulas using your computer, you may try to download a free program that does it. To install this program on your computer, go to the download page: <http://www.simulations-plus.com/software/medchem-designer/> and press a **Download** button.

This software is provided by a company that develops more sophisticated software for researchers and industry, so it is safe to download this program.

To complete downloading procedure, ask the questions about you. I myself just typed my home address and answered all questions pretending I am a school student. It worked. ©Mark Lukin



Figure 3: Pont du Gard aqueduct in Southern France. Built about 2000 years ago, it is still almost functional.

¹⁹Just to remind you. In this problem, you should use the Avogadro's law: if a mass of one molecule of some gas is X Da, then X *grams* of this gas will occupy a volume of 22.4 L at room temperature and atmospheric pressure.