## **MATH 10**

## ASSIGNMENT 2: OPEN AND CLOSED SETS

SEPT 24, 2017

**Definition 1.** A metric space is a set X with a distance function: for any  $x, y \in X$  we have a real number d(x, y) such that

- **1.** d(x,y) = d(y,x)
- **2.**  $d(x,y) \geq 0$  for any x,y
- **3.** d(x,y) = 0 if and only if x = y
- **4.** Triangle inequality:  $d(x,y) + d(y,z) \ge d(x,z)$ .

Usual examples are  $\mathbb{R}$ ,  $\mathbb{R}^2$ , ..., but there are other examples as well.

Given a point  $x \in X$  and a positive real number  $\varepsilon$ , we define  $\varepsilon$ -neighborhood of x by

$$B_{\varepsilon}(x) = \{ y \in X \mid d(x, y) < \varepsilon \}.$$

If  $S \subset X$ , denote by S' the complement of S. Then, for any  $x \in X$ , we can have one of three possibilities:

- 1. There is a neighborhood  $B_{\varepsilon}(x)$  which is completely inside S (in paritcular, this implies that  $x \in S$ ). Such points are called *interior points* of S; set of interior points is denoted by Int(S).
- **2.** There is a neighborhood  $B_{\varepsilon}(x)$  which is completely inside S' (in paritcular, this implies that  $x \in S'$ ). Thus,  $x \in \text{Int}(S')$ .
- **3.** Any neighborhood of x contains points from S and points from S' (in this case, we could have  $x \in S$  or  $x \in S'$ ). Set of such points is called the *boundary* of S and denoted  $\partial S$ .

**Definition 2.** A set S is called *open* if every point  $x \in S$  is an interior point: S = Int(S).

A set S is called closed if  $\partial S \subset S$ .

## Homework

1. Show that set  $\mathbb{R}^2$  with distance defined by

$$d((x_1, y_1), (x_2, y_2)) = |x_1 - x_2| + |y_1 - y_2|$$

is a metric space. (This distance is sometimes called Manhattan or taxicab distance — can you guess why?)

- **2.** For each of the following subsets of  $\mathbb{R}$ , find its interior and boundary and determine if it is open, closed, or neither.
  - (a) Set  $\mathbb{N} = \{1, 2, 3, \dots\}$ .
  - (b) Interval [0,1]
  - (c) Open interval (0,1)
  - (d) Interval [0,1).
  - (e) Set of all rational numbers
  - (f) Set consisting of just two points  $\{0, 1\}$
  - \*(g) Set  $x^3 + 2x + 1 > 0$

Are there any subsets of  $\mathbb{R}$  which are both open and closed?

- 3. Show that a set S is open if and only if its complement S' is closed.
- \*4. For a set S, let  $\overline{S} = S \cup \partial S = \{x \mid \text{ In any neighborhood of } x$ , there are elements of  $S\}$ . Prove that  $\overline{S}$  is closed. (It is called the closure of S.)
- 5. Show that union and intersection of two open sets is open. Is it true if we replace two sets by any collection of open sets?

Same question about closed sets.