

MATH 8: HANDOUT 14
EUCLIDEAN GEOMETRY 1: AXIOMS.

Euclidean geometry tries to describe geometric properties of various figures in the plane. Figures are understood as sets of **points**; we will use capital letters for points and write $P \in m$ for “point P lies in figure m ”, or “figure m contains point P ”. The notion of **point** can not be defined: it is so basic that it is impossible to explain it in terms of simpler notions. In addition, there are some other basic notions (**lines, distances, angle measures**) that can not be defined. Instead, we can state some basic properties of these objects; these basic properties are usually called **postulates** or **axioms of Euclidean geometry**. **All results in Euclidean geometry should be proven by deducing them from the axioms**; justifications “it is obvious”, “it is well-known”, or “it is clear from the figure” are not acceptable!

We allow use of all logical rules. We will also use all the usual properties of real numbers, equations, inequalities, etc.

For your enjoyment, take a look at the book which gave rise to Euclidean geometry and much more, Euclid’s *Elements*, dated about 300 BC, and used as the standard textbook for the next 2000 years. Nowadays it is available online at <http://math.clarku.edu/~djoyce/java/elements/toc.html>

1. BASIC OBJECTS

These objects are the basis of all our constructions: all objects we will be discussing will be defined in terms of these objects. No definition is given for these basic objects.

- Points (usually denoted by upper-case letters: A, B, \dots)
- Lines (usually denoted by lower-case letters: l, m, \dots)
- Distances: for any two points A, B , there is a non-negative number AB , called distance between A, B . The distance is zero if and only if points coincide.
- Angle measures: for any angle $\angle ABC$, there is a non-negative real number $m\angle ABC$, called the measure of this angle (more on this later).

We will also frequently use words “between” when describing relative position of points on a line (as in: A is between B and C) and “inside” (as in: point C is inside angle $\angle AOB$). We do not give full list of axioms for these notions; it is possible, but rather boring.

Having these basic notions, we can now define more objects. Namely, we can give definitions of

- interval, or line segment (notation: \overline{AB}): set of all points on line \overleftrightarrow{AB} which are between A and B , together with points A and B themselves;
- ray (notation: \overrightarrow{AB}): set of all points on the line \overleftrightarrow{AB} which are on the same side of A as B (Note that we have not defined the concept “on the same side” but will be using it in the future);
- angle (notation: $\angle AOD$): figure consisting of two rays with a common vertex;
- parallel lines: two distinct lines l, m are called parallel (notation: $l \parallel m$) if they do not intersect, i.e. have no common points. We also say that every line is parallel to itself (it is a rather convenient convention, which will make our lives easier – the intuition here is that parallel lines have the same “direction”).


2. FIRST AXIOMS

After we introduced some objects, including undefined ones, we need to have statements (*axioms*) that describe their properties. Of course, the lack of definition for undefined objects makes such properties impossible to prove. The goal here is to state the *minimal number* of such properties that we take for granted, just enough to be able to prove or derive harder and more complicated statements. Here are the first few axioms:


Axiom 1. For any two distinct points A, B , there is a unique line containing these points (this line is usually denoted \overleftrightarrow{AB}).

Axiom 2. If points A, B, C are on the same line, and B is between A and C , then $AC = AB + BC$

Axiom 3. If point B is inside angle $\angle AOC$, then $m\angle AOC = m\angle AOB + m\angle BOC$. Also, the measure of a straight angle is equal to 180° .



Axiom 4. Let line l intersect lines m, n and angles $\angle 1, \angle 2$ are as shown in the figure below (in this situation, such a pair of angles is called *alternate interior angles*). Then $m \parallel n$ if and only if $m\angle 1 = m\angle 2$.



In addition, we will assume that given a line l and a point A on it, for any positive real number d , there are exactly two points on l at distance d from A , on opposite sides of A , and similarly for angles: given a ray and angle measure, there are exactly two angles with that measure having that ray as one of the sides.

HOMEWORK

All problems below are important — please try to finish them all! Note that you may use all results that are presented in the previous sections. This means that you may use any theorem if you find it a useful logical step in your proof. The only exception is when you are explicitly asked to prove a given theorem, in which case you must understand how to draw the result of the theorem from previous theorems and axioms. Each step of your proof must be based on some previous, already proven, statement or an axiom.

1. It is important that you know some geometry notation.
 - (a) What does the symbol \parallel mean? How do you pronounce it? How would you read “ $a \parallel b$ ”?
 - (b) What does the symbol \perp mean? How would you say “ $a \perp b$ ”?
 - (c) Suppose you have two points X and Y . What is the difference between \overline{XY} , \overleftrightarrow{XY} , \overrightarrow{XY} ? What are each of these things called?
 - (d) Given three points E, F, G , what does $EF + FG$ mean?
 - (e) Given four points A, B, C, D , what does $m\angle ADC + m\angle BDC$ mean? If I tell you $m\angle ADC + m\angle BDC = 180^\circ$, does that tell you any information about $m\angle ADC$ or $m\angle BDC$?
 - (f) What does the symbol \triangle mean? For example, if A and B and C are points, what is $\triangle ABC$?
2.
 - (a) What is a proof? Give an example. Can you come up with an example that is not about geometry?
 - (b) What is an axiom? Give an example. Can you come up with an example that is not about geometry?
3. In this problem, you will make diagrams. Part of the purpose of this exercise is so that, when you think about geometry, the pictures in your notes or in your mind aren't all just the diagrams I draw out for you in class or on classwork sheets. You have to be able to draw or visualize configurations of lines other than the way they're set up in axiom 4, for example.

- (a) Given lines a, b, c , is it possible that $a \parallel b$ and $\neg(b \parallel c)$ but $a \parallel c$? Draw a diagram and then explain your reasoning on how to answer this question. (“explain” means, of course, in writing.)
- (b) Suppose we have parallel lines l, m . Let A, B, C be points on l , with B between A, C . Let X, Y, Z be points on m , with Y between X, Z . Is it possible for lines $\overleftrightarrow{AX}, \overleftrightarrow{BY}, \overleftrightarrow{CZ}$ to all intersect at one point? Draw a diagram of what this might look like.
- (c) Consider the diagram you drew in the previous part, with the lines l, m and the six points, and the three cross-lines that intersect at a point. Now consider the lines $\overleftrightarrow{AZ}, \overleftrightarrow{CX}$. Do these two lines intersect at a point on \overleftrightarrow{BY} ? Draw a diagram where this *is* the case, and then draw a second diagram where this *is not* the case.
- (d) Draw a rectangle that’s not a square, and draw it so that one of the bases is horizontal. Then draw one of the rectangle’s diagonals. Notice that, of the two right angles formed at the rectangle’s base, the rectangle’s diagonal splits one of those angles into two smaller angles. Which of the two angles is bigger - the one below the diagonal, or the one above the diagonal? Draw a second rectangle where the opposite relation holds true (for example, if the lower angle was bigger in your first rectangle, draw a second rectangle where the lower angle split by the diagonal is smaller).
4. Can you formulate Axiom 4 without referring to the picture (i.e. without using any statement such as “angles $\angle 1, \angle 2$ are as shown in the figure below”? You will have to introduce a number of points and have very clear notations.
5. The following logic and geometric statements come in equivalent pairs. Each logic statement has exactly one geometric statement that is equivalent to it. Match these statements into their equivalent pairs, with an explanation of why the pairs you chose are equivalent. [Note: the quantifier $\exists!$ stands for “there exists a unique...”, and \emptyset is an empty set.]

Geometric statements:

- (a) For any two distinct points there is a unique line containing these points.
- (b) Given a line and a point not on the line there exists a unique line through the given point that is parallel to the given line.
- (c) If two lines are parallel and another line intersects one of them, then it intersects the other one as well.
- (d) If two lines are parallel to the same line, then they are parallel to each other

Logic statements:

- (a) $\forall l \forall m$ such that $l \parallel m$ [$\forall n (n \cap l \neq \emptyset \rightarrow n \cap m \neq \emptyset)$]
- (b) $\forall A \forall B$ such that $A \neq B$ [$\exists! l (A \in l \wedge B \in l)$]
- (c) $\forall l \forall m$ [$(\exists n$ such that $n \parallel l \wedge n \parallel m) \rightarrow (l \parallel m)$]
- (d) $\forall l \forall A$ such that $A \notin l$ [$\exists! m (A \in m \wedge m \parallel l)$]