

MATH 10
ASSIGNMENT 25: PUBLIC KEY CRYPTOGRAPHY
APR 30, 2023

SUMMARY OF PREVIOUS RESULTS

Theorem 1. *If a is relatively prime to n , then $a^{\varphi(n)} \equiv 1 \pmod{n}$, where $\varphi(n)$ is the Euler's function:*

$\varphi(n)$ = number of remainders modulo n which are relatively prime to n .

In particular, for prime p , we have $a^{p-1} \equiv 1 \pmod{p}$ for any a not divisible by p .

Chinese remainder theorem also gives the following result:

Theorem 2. *If m, n relatively prime, then $\varphi(mn) = \varphi(m)\varphi(n)$.*

PUBLIC KEY CRYPTOGRAPHY

In RSA model of public key cryptography, the encoding and decoding is done as follows:

First, a person (call him Bob) chooses a number n which should be a product of two primes $n = pq$. In real life, p, q are chosen so that each is about 100 digits long; obviously, in our examples we will use smaller numbers. He also chooses two numbers d, e such that

$$de \equiv 1 \pmod{\varphi(n)}$$

where $\varphi(n)$ is Euler's function. Pair (e, n) is called the public key; pair (d, n) is called the private key. The public key is made widely known (posted online); the private key (and the numbers p, q) he keeps to himself.

Now if someone (say, Alice) wants to send Bob an encrypted email, it can be done as follows. First, the text message is broken into blocks of fixed length, and each block is converted to a number (call it P) between 1 and n . Next, each such number is replaced by an encrypted number C defined by

$$C = P^e \pmod{n}$$

This sequence of encrypted numbers is sent in an email to Bob.

To decode this, Bob uses his private key: he recovers P from C by using the formula $P = C^d \pmod{n}$ (since $C^d \equiv P^{de} \equiv P^1 \equiv P \pmod{n}$).

The remarkable thing is that here, different keys are used for encrypting and decrypting. What is more, the only way to recover d from e requires finding $\varphi(n)$ — which in turn requires finding the prime factors p, q . Since n is about 200 digit long, factoring it is extremely hard: at the moment, even the most powerful computers can not do it in reasonable time. Thus, for all practical purposes, even if you know the public key (e, n) , you can not find the private key (d, n) . Thus, anyone can use the public key to encrypt a message to Bob — but only Bob would be able to read it.

You can find the demos here: <http://cisnet.baruch.cuny.edu/hollowczak/classes/9444/rsademo/> and here <http://www.cs.pitt.edu/~kirk/cs1501/notes/rsademo/index.html>

This scheme has also many other applications, some of which were discussed in class.

- *1. Chinese Remainder Theorem claims that if m, n are relatively prime, then any system of congruences

$$\begin{aligned}x &\equiv a_1 \pmod{m} \\x &\equiv a_2 \pmod{n}\end{aligned}$$

has a solution, and the solution is unique up to adding a multiple of mn . In other words, given a pair of remainders mod m and mod n , we can uniquely construct a remainder mod mn .

- (a) Deduce from this that there exists a bijection $\mathbb{Z}_{mn} \rightarrow \mathbb{Z}_m \times \mathbb{Z}_n$ which preserves addition and multiplication.
- (b) Prove that $\varphi(mn) = \varphi(m)\varphi(n)$ (as before, assuming that m, n are relatively prime).
2. Use results of the previous problem to write a general formula for $\varphi(n)$, where $n = p_1^{k_1} \dots p_n^{k_n}$. Find $\varphi(15)$; $\varphi(100)$; $\varphi(1001)$; $\varphi(240)$; $\varphi(30000)$; $\varphi(96)$.
3. Compute the remainder $2^{2170} \pmod{1001}$. (Hint: do you remember the factorization of 1001?)
4. Compute the last two digits of 2011^{970} .
5. (a) Find the inverse of 17 mod 24, i.e. solve $17e \equiv 1 \pmod{24}$.
(b) If a is relatively prime to 35, and $b \equiv a^{17} \pmod{35}$, then one can write $a \equiv b^e \pmod{35}$ for $e = ?$.
6. Assume that a person uses RSA encoding, with public key ($e = 13, n = 77$), so the encoding is done by the formula

$$C = P^{13} \pmod{77}$$

Can you find the key that should be used for decoding?

7. Assume that a person uses RSA encoding, with public key ($e = 7, n = 437$). Break the encoding, by factoring n and finding d . Use it to decode the following secret message:

$$426 \ 4 \quad 215 \ 1 \ 215$$

8. Two people want to play “stone, scissors, paper” online. However, none can trust the other. They could play by mail using the following trick: each writes his answer on a piece of paper, locks it in a strong box, and sends to his partner — but without the key. Only after both receive the boxes, they send each other the key.
Can you think of a similar trick which would allow them to play online, using public key cryptography?