

November 13, 2022

Algebra.

Solutions to some homework problems.

1. **Problem.** Write the first few terms in the following sequence ($n \geq 1$),

$$n \text{ fractions } \left\{ \begin{array}{l} \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \dots}}} \\ \dots + \frac{1}{1+x} \end{array} \right. = f_n$$

- a. Try guessing the general formula of this fraction for any n .
- b. Using mathematical induction, try proving the formula you guessed.

Solution. $n = 1: f_1 = \frac{1}{1+x}; n = 2: f_2 = \frac{1}{1 + \frac{1}{1+x}} = \frac{1+x}{2+x}; n = 3, f_3 = \frac{1}{1 + \frac{1}{1 + \frac{1}{1+x}}} = \frac{2+x}{3+2x}; n = 4, f_4 = \frac{1}{1 + \frac{1}{1 + \frac{1}{1 + \frac{1}{1+x}}}} = \frac{3+2x}{5+3x}; f_5 = \frac{5+3x}{8+5x}; \dots$

From the definition, we can write the recurrence, $f_{n+1} = \frac{1}{1+f_n}$. We note, that if $f_n = \frac{a_n+b_nx}{c_n+d_nx}$, then $f_{n+1} = \frac{c_n+d_nx}{(a_n+c_n)+(b_n+d_n)x}$. Hence, in each next term, f_{n+1} , in the sequence, the numerator is equal to the denominator of the previous term, f_n , while the numbers in the denominator are the sums of the corresponding numbers in the numerator and the denominator of the previous term, f_n , thus forming the Fibonacci sequence, $\{F_n\} = \{1,1,2,3,5,8,13, \dots\}$. We can thus guess,

- a. n fractions: $f_1 = \frac{1}{1+x}, f_n = \frac{F_n+F_{n-1}x}{F_{n+1}+F_nx}, n > 1$
- b. Base: $f_2 = \frac{1+x}{1+2x}$

Induction: Using the recurrence implied in the definition,

$$f_{n+1} = \frac{1}{1+f_n} = \frac{1}{1 + \frac{F_n+F_{n-1}x}{F_{n+1}+F_nx}} = \frac{F_{n+1}+F_nx}{F_{n+1}+F_n+F_nx+F_{n-1}x} = \frac{F_{n+1}+F_nx}{F_{n+2}+F_{n+1}x}$$

2. **Problem.** Can you prove that,

a.

$$\frac{3+\sqrt{17}}{2} = 3 + \frac{2}{3 + \frac{2}{3 + \frac{2}{\dots}}} ?$$

b. $1 = 3 - \frac{2}{3 - \frac{2}{3 - \frac{2}{\dots}}} ?$

c.

$$\frac{4}{2 + \frac{4}{2 + \frac{4}{\dots}}} = 1 + \frac{1}{4 + \frac{1}{4 + \frac{1}{\dots}}} ?$$

Find these numbers?

Solution. Consider a general continued fraction,

$$x = a + \frac{b}{a + \frac{b}{a + \frac{b}{\dots}}}$$

If a number exists, which is equal to the above infinite continued fraction, then it must satisfy the equation, $x = a + \frac{b}{x} \Leftrightarrow x^2 - ax - b = 0$

$\Leftrightarrow x = \frac{a}{2} \pm \sqrt{\left(\frac{a}{2}\right)^2 + b}$. If a and b are positive, then x must also be

positive, so $x = \frac{a}{2} + \sqrt{\left(\frac{a}{2}\right)^2 + b}$.

a. Following the above argument with $a = 3$, $b = 2$, we obtain, $x = \frac{3}{2} +$

$$\sqrt{\left(\frac{3}{2}\right)^2 + 2} = \frac{3+\sqrt{17}}{2}$$

b. In this case, $a = 3$, but $b = -2$ is negative. Applying the above considerations naively, we obtain, $x = 3 - \frac{2}{x} \Leftrightarrow x^2 - 3x + 2 = 0$
 $\Leftrightarrow (x - 1)(x - 2) = 0$, i.e. there are two equally "legitimate" answers, $x = 1$, or $x = 2$. What this means, is that assumption that there exist unique number encoded by the given infinite continued fraction is wrong: there exists no such number! In fact, this can also be understood by looking at finite truncations approximating this

continued fraction. If the continued fraction is truncated after subtracting 2 and before division by 3, then it is equal to 1,

$$3 - \frac{2}{3-2} = 1, 3 - \frac{2}{3 - \frac{2}{3-2}} = 1, \dots$$

If, on the other hand, the truncation is after division by 3 and before subtracting 2, then we obtain a sequence of numbers approaching 2,

$$3 - \frac{2}{3} = 2\frac{1}{3}, 3 - \frac{2}{3 - \frac{2}{3}} = 2\frac{1}{7}, 3 - \frac{2}{3 - \frac{2}{3 - \frac{2}{3}}} = 2\frac{1}{15}, \dots$$

c. Denote

$$x = \frac{4}{2 + \frac{4}{2 + \frac{4}{2 + \dots}}} = \frac{4}{2 + x}$$

Then, $x^2 + 2x - 4 = 0 \Leftrightarrow x = -1 \pm \frac{\sqrt{5}}{2}$, and $x > 0$. Hence, $x = -1 + \frac{\sqrt{5}}{2}$.

Similarly, denote

$$y = \frac{1}{4 + \frac{1}{4 + \frac{1}{4 + \dots}}} = \frac{1}{4 + y}$$

Then, $y^2 + 4y - 1 = 0 \Leftrightarrow y = -2 \pm \frac{\sqrt{5}}{2}$, and $y > 0$. Hence, $y = -2 + \frac{\sqrt{5}}{2}$, and $1 + y = -1 + \frac{\sqrt{5}}{2} = x$.